

Naturetrek Bargain Birds Tour

South Africa

9th – 16th February 2010



Bateleur, seen on the last morning

Tour Summary

Tour Participants:

Duncan & Janet Fraser
Susan Luce
Paul Marchant
Anthony O'Mahony
Colin & Pauline Raymond
Edward & Penelope Raynor
Richard & Penelope Scriven
Ann Yates

Tour Leader: Errol de Beer, assisted by John Davies

Photo Acknowledgments: Errol de Beer

Total Distance Travelled: 1743 km

Temperature Range: 13° – 39° Celsius

Altitude Variation: 260 – 2100 meters above sea level

Total Number of Birds Seen: 318

Total Number of Mammals Seen: 39

Trip Report Compiled By: Errol de Beer

Note:

The species mentioned in the report are only some of the species seen at each locality. Please refer to detailed checklists at end of report as to all species seen on this specific tour.

Trip Breakdown

Day 1: Tuesday, 9th February 2010 ~ Dullstroom

Route: OR Tambo International Airport to Dullstroom.

Distance: 294 km.

Weather: Clear.

Day 2: Wednesday, 10th February 2010 ~ Dullstroom

Route: Dullstroom area.

Distance: 118 km.

Weather: Hot and clear.

Day 3: Thursday, 11th February 2010 ~ Blyde River Canyon

Route: Dullstroom, Lydenburg, Mount Sheba, Pilgrim's Rest, Graskop to Blyde Aventura.

Distance: 212 km.

Weather: Clear for most of the day with some cloud in the afternoon.

Day 4: Friday, 12th February 2010 ~ Satara, Kruger National Park

Route: Blyde Canyon, Abel Erasmus Pass, Strijdom Tunnel, Orpen Gate to Satara Rest Camp.

Distance: 205 km.

Weather: Clear and very hot.

Day 5: Saturday, 13th February 2010 ~ Satara, Kruger National Park

Route: Satara region.

Distance: 94 km.

Weather: Clear and very hot.

Day 6: Sunday, 14th February 2010 ~ Skukuza, Kruger National Park

Route: Satara to Skukuza via Tshokwane Picnic Site.

Distance: 112 km.

Weather: Partly cloudy and hot with rain in the afternoon.

Day 7: Monday, 15th February 2010 ~ Pretoriuskop, Kruger National Park

Route: Skukuza to Pretoriuskop.

Distance: 94 km.

Weather: Partly cloudy and hot.

Day 8: Tuesday, 16th February 2010 ~ Departure

Route: Pretoriuskop to Johannesburg

Distance: 523 km.

Weather: Partly cloudy and hot.

TRIP REPORT

Day 1: Tuesday, 9th February 2010 ~ Dullstroom

The late arrival of the flight from London saw us off to a slightly later start than initially anticipated but the beautiful weather boded well for the rest of the day. We made our first stop at a small pan (pond) along the N12 not too far out of the city. The pan was teeming with life and we soon ticked Fulvous Duck, Cape Shoveler, Southern Pochard and Red-billed Teal. Whiskered Terns were actively hunting over the pan and we were further entertained by the striking Red and Yellow-crowned Bishops. Roadside raptors included numerous Amur Falcon as well as the odd Steppe Buzzard. Upon our arrival in Dullstroom we decided to give everyone a chance to settle in before heading off for some afternoon birding. Our first bird of the afternoon was African Olive-Pigeon soon followed by Alpine Swift and Speckled Mousebird. From town we headed down to the Municipal Dams where our main target was the rather rare Cape Eagle-Owl. Along the entrance road we were all treated to good views of a Long-crested Eagle soaring low over the exotic plantations. Birding around the dams was rather quiet with only a few Brown-throated Martins flying low over the water. We decided to skirt around the base of the hill rather than climb all the way up, which left us with only a very small climb to get to a decent vantage point from where we could scan for the Owl. Along here we had splendid views of Dark-capped Yellow-Warbler and a male Malachite Sunbird in all its glory. The Owl proved rather elusive and although we could hear it all the time we only managed to locate it just before dark, nevertheless we managed great scope views of this superb bird.

Daily Total: 82

Trip Total: 82

Mammal total for the day: 4

Bird of the day: Cape Eagle-Owl

Day 2: Wednesday, 10th February 2010 ~ Dullstroom



We started the day at 05h30 and had coffee and rusks a little way down the "De Berg" road. We stopped alongside a small wetland and it was barely seconds after stopping that we heard Red-chested Flufftail calling. We tried in vain to entice it out of cover and had to be satisfied with the call only. One of the highlights of the day was crippling views of a Common Cuckoo hunting caterpillars from a small dry bush on a hillside, the bird was very confiding and allowed us enough time to have superb scope views. Cape Canaries were common as usual and further along we found Buff-streaked Chat and Jackal Buzzard. Good birding did not allow us to advance at any great pace and by the time we reached the Verloren Vallei Nature Reserve Turn-off it was almost time to head back for breakfast. A Cape Robin-Chat was seen briefly and just before turning back we all had good views of a lone Oribi, a rather rare antelope, so quite a nice addition to our mammal list.

After a hearty breakfast a small group of us decided to explore the Kruisfontein road while the others opted for a midday break. It was very hot and birding was really slow but we managed cracking views of Red-winged Francolin, usually quite an unobtrusive species that is more readily heard than seen. Other good sightings during this stint were Drakensberg Prinia and a juvenile Rock Kestrel.

At 14h00 we joined up with the rest of the group and headed back to Verloren Vallei. Just past Linga Longa (a lodge near the nature reserve) we encountered a pair of Blue Crane in a recently ploughed field. They were joined by a good number of Southern Bald Ibis. Reports from a passerby alerted us to the presence of Wattled Cranes on the high ground which had us scrambling to where they were seen, but to no avail, a thorough search of the area proved negative. During this search we did however come up trumps with both Secretarybird and Denham's Bustard. The high ground had more good birds instore for us and Yellow-breasted Pipit and Eastern Long-billed Lark were both ace birds. We reached the main tarred road and the Protea's (known locally as Suikerbos, *Protea caffra*) here held significant numbers of Gurney's Sugarbird.

After an amazing dinner we all retired for a well deserved rest.

Daily Total: 88
New Birds: 39
Trip Total: 127
Mammal total for the day: 8
Bird of the day: Blue Crane

Day 3: Thursday, 11th February 2010 ~ Blyde River Canyon

Our journey saw us off to an early start from Dullstroom on our way to Mount Sheba. Common roadside birds included White Stork and Black-shouldered Kite. Upon our arrival at the turn-off to Mount Sheba we were greeted by a beautiful pair of Secretarybirds right next to the road, improving our previous day's sighting. A Red-winged Francolin poked its head out of the grass but resisted all John efforts to flush it. A few short stops before reaching the first patch of Afro-montane Forest produced good views of Cape Grassbird and Bar-throated Apalis. From here we made a steady push down to the hotel for breakfast.

Shortly after breakfast we found a male Southern Double-collared Sunbird feeding in a bottlebrush tree right outside reception. From here we made our way into the forest and along the first path we managed to startle a couple of juvenile Black Sparrowhawks, and this was followed by good sightings of Yellow-throated Woodland Warbler. Deeper into the forest we found Cape Batis, Chorister Robin-Chat, Knysna Turaco, Yellow-streaked Greenbul and last but not least, the most amazing views of Narina Trogon. A pair of Klipspringer finished off a great morning at Mt. Sheba.

We stopped for lunch at Harry's Pancakes in Graskop followed by a brief stop at a patch of flowers where we recorded a number of sunbird species which included amongst others Amethyst, Greater Double-collared and Malachite Sunbirds. Upon our arrival at the Blyde Resort we headed to the top view point from where we could see the Blyde Canyon in all its glory. We were most surprised to have two different sighting of Taita Falcon from here.

Daily Total: 91
New Birds: 25
Trip Total: 146
Mammal total for the day: 7
Bird of the day: Narina Trogon

Day 4: Friday, 12th February 2010 ~ Satara, Kruger National Park

Early morning coffee was followed by a walk down to the lower viewpoint where we recorded numerous birds, including Mocking Cliff-Chat, Cinnamon-breasted Bunting, Yellow-breasted Apalis, Rock Martin, Red-capped Robin-Chat, African Paradise Flycatcher and Striped Pipit. Mountain Wagtail and White-throated Robin-Chat were recorded at a small dam near the camping ground, and not far from there a Streaky-headed Seed-Eater.

After breakfast we packed up and headed for the Abel Erasmus Pass, where on our arrival we were greeted by a smiling Michael (the local guide) who was ready to show us the diminutive Taita Falcon perched on a small ledge on the cliff face. This is usually a real highlight of the trip but probably less so today since we already had good views of them hunting the previous day. We saw our first Cape Vultures of the trip here and Red-

winged Starlings were everywhere. Some of the group had a possible African Cuckoo Hawk sighting but we were too late to confirm the id. From here it was pretty much a straight drive to Orpen gate, our entry point into the world famous Kruger National Park. There was off-course the compulsory lunch stop just outside the town of Klaserie.

It was extremely hot when we arrived at the gate and birding proved quite slow. Once inside the park we ticked new birds at regular intervals , good birds included Long-tailed Paradise Whydah, Jameson's Firefinch, Long-billed Crombec, Magpie Shrike and Burchell's Starling. Further along we had good sightings of Bateleur and White-headed Vulture, perched in the same tree. By now we were pushing a bit harder in order to reach the camp gates before closing time, but as we drove along I saw two figures in the road and immediately shouted "cats"!! On closer inspection they proved to be Leopard. What a sighting and as if that was not enough we bumped into another Leopard barely three kilometres from camp. Needless to say we were about ten minutes late, but fortunately the camp staff at the gate were friendly and waved us through with a smile.

Daily Total: 122
New Birds: 65
Trip Total: 211
Mammal total for the day: 14
Bird of the day: Taita Falcon

Day 5: Saturday, 13th February 2010 ~ Satara, Kruger National Park

We started off just when the camp gates opened at 05h30, heading east down the S100 towards Nwanetsi Picnic spot. Just outside the gates we got our first new bird of the day in the form of a Burnt-necked Eremomela, followed shortly thereafter by a Harrier of some sorts, unfortunately the light was too bad and the bird in a hurry somewhere so no conclusive ID was made. As if we had not been lucky enough seeing three Leopards the previous evening another prime male sauntered across the road, I quickly radioed John to get back and fortunately they made it just in time before the cat disappeared into the tall grass. In the same area we managed to record Yellow-bellied Eremomela, a bird not all that often recorded on birding trips and Golden-breasted Bunting. A Giant Kingfisher was seen perched along the banks of the Nwanetsi River and we heard and located numerous Brown-headed Parrots. Probably one of the best sightings of the day was that of a extremely confiding Olive-tree Warbler, and Ed explained to us that the downward tail pumping action is highly distinctive for this species and this was witnessed firsthand by all. The air was filled with the song of Monotonous Lark along the S41 and we managed to get superb views of this Lark, which, when not singing, can be rather unobtrusive. A quick stop at the picnic spot turned up Bushbuck down in the gorge and from there we headed to Sweni hide where we recorded Malachite Kingfisher, Black Crake, Common Moorhen, Hamerkop and Black-crowned Night-Heron. A slow drive back to camp via the H6 delivered our first African Buffalo, an old Dagga Boy, at Shishangani waterhole.

Our afternoon drive to Nsemani dam produced hordes of Marabou , Yellow-billed and Woolly-necked Storks, and from there a short drive to Girivana waterhole proved worthwhile as we spotted out first of four separate Dwarf Bittern sightings here.

Daily Total: 133
New Birds: 55
Trip Total: 266
Mammal total for the day: 18
Bird of the day: Dwarf Bittern



Day 6: Sunday, 14th February 2010 ~ Skukuza, Kruger National Park

The mornings birding started off rather slow but a drive down to Sweni waterhole produced our first Spotted Hyena for the trip but unfortunately no Lions. Shortly afterwards we came upon a huge Red-billed Quelea colony and in close association we found a number of Lesser Spotted Eagle, obviously feeding on the rich pickings here. A short detour to Nkaya Pan was rather uneventfull other than for Lesser Masked Weaver and Brown-crowned Tchagra.

A short stop at an unmarked, inundated wetland saw us come away with another Dwarf Bittern, Little Bittern and Red-billed Firefinch. Other good birds along this stretch before arriving at Kumana Dam included Martial Eagle, Desert Cisticola, Tawny Eagle, Wahlberg's Eagle, White-backed Vulture, Gabar Goshawk, Orange-breasted Bush-Shrike and Marsh Warbler. At Kumana dam we watched as a huge herd of African Buffalo came down to drink, a careful scan through the Oxpeckers revealed only Red-billed Oxpecker but the buffalo did flush a Red-billed Teal, a rather rare bird in the park. There were quite a number of Comb Duck around with a lone Black-winged Stilt and a small number of Water Thick-knee completing the line-up.



We arrived at Tshokwane Picnic Spot just in time for lunch. Greater Blue-eared Starlings and African Mourning Dove are somewhat tame here. They regularly jump onto the tables in search of a scrap of food. After lunch we birded the grounds, and here we had good looks at African Green Pigeon, Southern Boubou, Eurasian Golden Oriole and Village Indigobird. Two species of Woodpecker in close proximity gave everyone the chance to compare the differences, spotted breast and white throat for Bennet's Woodpecker and streaked breast and throat for Golden-tailed Woodpecker. The find of the day however came in the form of a European Nightjar in the tree we parked the vehicles under, great stuff!!

On the way down to Skukuza we saw both Hooded and Lappet-faced Vultures and along the Sabie River we ticked Jacobin Cuckoo, Southern Carmine Bee-eater, White-fronted Bee-eater, Black-crowned Tchagra and Jacobin Cuckoo. This stretch hosted numerous herds of Elephant as well as one old tusker, truly a magnificent animal.

On reaching Skukuza some of the group opted for a sunset drive aboard one of the open safari vehicles belonging to the park, while the rest of us took a well deserved rest. They managed to find a number of White Rhino and Spotted Eagle-Owl on their excursion. On their return we headed to the Selati restaurant for a splendid dinner.

Daily Total: 139
New Birds: 23
Trip Total: 289
Mammal total for the day: 21
Bird of the day: European Nightjar

Day 7: Monday, 15th February 2010 ~ Pretoriuskop, Kruger National Park

This morning we decided to have a quick walk around camp before heading out to the Skukuza Golf Course where we would have our breakfast at the clubhouse. In camp we found a number of our target birds with some of the more noteworthy sightings being Purple-crested Turaco, Terrestrial Brownbul, Sombre Greenbul, Ashy Flycatcher, Collared Sunbird and White-browed Robin-Chat. Then, with bags packed we headed for the golf course, finding Senegal Lapwing en-route. At the golf course a short walk produced good sightings of numerous Little Bittern, Lesser Swamp Warbler and Grey Tit-Flycatcher. Breakfast at the clubhouse was definitely the best we had in the park, and while watching the Hippo's in the Lake. After breakfast we took a stroll through the nursery where we ticked Scarlet-chested Sunbird before heading for the Lake Panic Bird Hide, a favourite spot amongst visiting bird watchers. Fortunately the hide was fairly empty, but not much was happening. We did however find African Darter, Green-backed Heron and African Jacana. On the way out we found two immature African Goshawks, probably the same birds that I saw here in January.

We still had a fair distance to cover to our next camp, Pretoriuskop, and we proceeded with a few stops along the way, most notably for a White Rhino mother with a calf at Napi Boulder. At Pretoriuskop we checked in and had lunch before going on an afternoon drive along Fayi Loop that proved very productive, with good birds such as Green-capped Eremomela, Common Scimitarbill, Striped Kingfisher and Dark-Chanting Goshawk.

On our way to dinner at Pretoriuskop this evening we had cracking views of a Pearl-spotted Owlet.

Daily Total: 135
New Birds: 17
Trip Total: 306
Mammal total for the day: 20
Bird of the day: Senegal Lapwing

Day 8: Tuesday, 16th February 2010 ~ Departure

Our final morning started with a bit of birding in the camp before breakfast. Interesting birds included Black Cuckooshrike, White-bellied Sunbird, Willow Warbler and Cape Glossy Starling. At breakfast we got word of two Lions, a male and female, near Shabeni hill and since this was the only one of the Big 5 still outstanding we decided to go and check it out. On the way there we managed to flush an African Crake from the roadside near a small stream and we also managed good looks at Croaking Cisticola here. We drove up the S7 and just as we passed Shabeni hill we saw two vehicles parked and right next to them there were two Lions laying in the road. We pulled up slowly and watched these awesome animals for a good while.

From here we headed out of the park with a short stop at the Lowveld Botanical Gardens in Nelspruit for lunch. We did manage one new bird in the gardens in the form of a Tambourine Dove, but unfortunately it was not seen by all. After a really delicious lunch we tackled the long journey to Johannesburg with one more critical stop in mind, Wonderfontein Pans. This stop never fails in adding at least a couple of new birds and today was no exception. As we stopped, John spotted the first new one, a White-backed Duck followed by Maccoa Duck and Great Crested Grebe. White-winged and Whiskered Terns were also present



and were quartering low over the water. The pan on the opposite side of the road accounted for a few more new birds. These included African Spoonbill, Greater Flamingo, Cape Teal, Common Greenshank and Ruff. Our last new bird to make the list was South African Cliff-Swallow at a bridge over the Bronkhorst Spruit (Spruit, being Afrikaans for stream). At the airport we said our final farewells and wished everyone a safe journey home.

Daily Total: 100
New Birds: 12
Trip Total: 319
Mammal total for the day: 7
Bird of the day: African Crake

Summary

Total Bird Species: 319; Total endemics / near endemic birds: 36; Total Mammal species: 39

What an incredible trip. My first trip for Naturetrek and hopefully not the last. There were 99 recorded mammal sightings and 890 recorded bird sightings for the tour, which gives you some idea of the numbers of birds and animals seen on a daily basis. Despite the extremely hot temperatures we succeeded in finding an average of 111 birds per day and some rather good ones at that, combined with great sightings of the Big 5 mammals. This was an amazingly successful trip.

Top ten birds for the tour: Blue Crane; Cape Eagle-Owl; Narina Trogon; Red-winged Francolin; Taita Falcon; Senegal Lapwing; African Crake; Dwarf Bittern; European Nightjar; Lappet-faced Vulture.

Bird List (E = Endemic, NE = Near Endemic)

| Common name | Scientific name | Total | Day | | | | | | | |
|--------------------------------|---------------------------------|-------|-----|---|---|---|---|---|---|---|
| | | | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 |
| Great Crested Grebe | <i>Podiceps cristatus</i> | 2 | X | | | | | | | X |
| Little Grebe | <i>Tachybaptus ruficollis</i> | 3 | X | | | | | | X | X |
| White-breasted Cormorant | <i>Phalacrocorax lucidus</i> | 3 | X | X | | | | | | X |
| Reed Cormorant | <i>Phalacrocorax africanus</i> | 3 | X | X | | | | | | X |
| African Darter | <i>Anhinga rufa</i> | 2 | X | | | | | | | X |
| Grey Heron | <i>Ardea cinerea</i> | 5 | X | | | | | X | X | X |
| Black-headed Heron | <i>Ardea melanocephala</i> | 6 | X | X | X | X | | | X | X |
| Great Egret | <i>Casmerodius albus</i> | 1 | X | | | | | | | |
| Yellow-billed Egret | <i>Egretta intermedia</i> | 1 | X | | | | | | | |
| Cattle Egret | <i>Bubulcus ibis</i> | 6 | X | X | X | X | X | | | X |
| Green-backed Heron | <i>Butorides striatus</i> | 3 | | | | | X | X | X | |
| Black-crowned Night-Heron | <i>Nycticorax nycticorax</i> | 1 | | | | | X | | | |
| Little Bittern | <i>Ixobrychus minutus</i> | 2 | | | | | | X | X | |
| Dwarf Bittern | <i>Ixobrychus sturmii</i> | 2 | | | | | X | X | | |
| Hamerkop | <i>Scopus umbretta</i> | 6 | | X | X | | X | X | X | X |
| White Stork | <i>Ciconia ciconia</i> | 7 | X | X | X | X | X | X | | X |
| Black Stork | <i>Ciconia nigra</i> | 1 | | | | | X | | | |
| Woolly-necked Stork | <i>Ciconia episcopus</i> | 1 | | | | | X | | | |
| African Openbill | <i>Anastomus lamelligerus</i> | 1 | | | | | X | | | |
| Marabou Stork | <i>Leptoptilos crumeniferus</i> | 4 | | | | X | X | X | X | |
| Yellow-billed Stork | <i>Mycteria ibis</i> | 2 | | | | | X | X | | |
| African Sacred Ibis | <i>Threskiornis aethiopicus</i> | 3 | X | | X | | | | | X |
| Southern Bald Ibis (E) | <i>Geronticus calvus</i> | 1 | | X | | | | | | |
| Glossy Ibis | <i>Plegadis falcinellus</i> | 2 | X | | | | | | | X |
| Hadeda Ibis | <i>Bostrychia hagedash</i> | 8 | X | X | X | X | X | X | X | X |
| African Spoonbill | <i>Platalea alba</i> | 1 | | | | | | | | X |
| Greater Flamingo | <i>Phoenicopterus ruber</i> | 1 | | | | | | | | X |
| White-faced Duck | <i>Dendrocygna viduata</i> | 3 | | | | | X | X | X | |
| Fulvous Duck | <i>Dendrocygna bicolor</i> | 1 | X | | | | | | | |
| White-backed Duck | <i>Thalassornis leuconotus</i> | 1 | | | | | | | | X |
| Egyptian Goose | <i>Alopochen aegyptiaca</i> | 7 | X | X | X | | X | X | X | X |
| Yellow-billed Duck | <i>Anas undulata</i> | 4 | X | X | X | | | | | X |
| Cape Teal | <i>Anas capensis</i> | 1 | | | | | | | | X |
| Red-billed Teal | <i>Anas erythrorhyncha</i> | 2 | X | | | | | X | | |
| Cape Shoveler (E) | <i>Anas smithii</i> | 2 | X | | | | | | | X |
| Southern Pochard | <i>Netta erythrophthalma</i> | 1 | X | | | | | | | |
| Comb Duck | <i>Sarkidiomis melanotos</i> | 3 | | | | | X | X | | X |
| Spur-winged Goose | <i>Plectropterus gambensis</i> | 3 | | X | X | | | | | X |
| Maccoa Duck | <i>Oxyura maccoa</i> | 1 | | | | | | | | X |
| Secretarybird | <i>Sagittarius serpentarius</i> | 2 | | X | X | | | | | |
| Hooded Vulture | <i>Necrosyrtes monachus</i> | 2 | | | | | X | X | | |
| Cape Vulture (E) | <i>Gyps coprotheres</i> | 2 | | | | X | | X | | |
| (African) White-backed Vulture | <i>Gyps africanus</i> | 5 | | | | X | X | X | X | X |
| Lappet-faced Vulture | <i>Torgos tracheliotus</i> | 1 | | | | | | X | | |
| White-headed Vulture | <i>Trigonoceps occipitalis</i> | 1 | | | | X | | | | |

| Common name | Scientific name | Total | Day | | | | | | | |
|---------------------------|---------------------------------|-------|-----|---|---|---|---|---|---|---|
| | | | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 |
| Yellow-billed Kite | <i>Milvus aegyptius</i> | 6 | X | | | X | X | X | X | X |
| Black-shouldered Kite | <i>Elanus caeruleus</i> | 8 | X | X | X | X | X | X | X | X |
| Tawny Eagle | <i>Aquila rapax</i> | 4 | | | | X | X | X | X | |
| Lesser Spotted Eagle | <i>Aquila pomarina</i> | 2 | | | | | X | X | | |
| Wahlberg's Eagle | <i>Aquila wahlbergi</i> | 5 | | | | X | X | X | X | X |
| Long-crested Eagle | <i>Lophaetus occipitalis</i> | 3 | X | | X | | | | | X |
| Martial Eagle | <i>Polemaetus bellicosus</i> | 2 | | | | | X | X | | |
| Brown Snake-Eagle | <i>Circaetus cinereus</i> | 3 | | | | X | | X | X | |
| Black-chested Snake-Eagle | <i>Circaetus gallicus</i> | 2 | | | | X | | | X | |
| Bateleur | <i>Terathopius ecaudatus</i> | 5 | | | | X | X | X | X | X |
| African Fish Eagle | <i>Haliaeetus vocifer</i> | 3 | | | | X | X | X | | |
| Steppe Buzzard | <i>Buteo buteo</i> | 5 | X | X | X | X | | | | X |
| Jackal Buzzard (E) | <i>Buteo rufofuscus</i> | 3 | | X | X | X | | | | |
| Black Sparrowhawk | <i>Accipiter melanoleucus</i> | 1 | | | X | | | | | |
| African Goshawk | <i>Accipiter tachiro</i> | 1 | | | | | | | X | |
| Gabar Goshawk | <i>Micronisus gabar</i> | 5 | X | | | X | X | X | X | |
| Dark Chanting Goshawk | <i>Melierax metabates</i> | 1 | | | | | | | X | |
| Peregrine Falcon | <i>Falco peregrinus</i> | 3 | | | X | X | X | | | |
| Eurasian Hobby | <i>Falco subbuteo</i> | 2 | X | | | | | | X | |
| Taita Falcon | <i>Falco fasciinucha</i> | 2 | | | X | X | | | | |
| Amur Falcon | <i>Falco amurensis</i> | 6 | X | X | X | X | | X | | X |
| Rock Kestrel | <i>Falco tinnunculus</i> | 2 | | X | X | | | | | |
| Crested Francolin | <i>Peliperdix sephaena</i> | 3 | | | | X | X | X | | |
| Red-winged Francolin | <i>Scleroptila levaillantii</i> | 2 | | X | X | | | | | |
| Natal Francolin (NE) | <i>Pternistes natalensis</i> | 5 | | | | X | X | X | X | X |
| Swainson's Spurfowl (NE) | <i>Pternistes swainsonii</i> | 4 | | X | | X | X | X | | |
| Helmeted Guineafowl | <i>Numida meleagris</i> | 8 | X | X | X | X | X | X | X | X |
| Blue Crane (E) | <i>Anthropoides paradiseus</i> | 1 | | X | | | | | | |
| African Crane | <i>Crecoptis egregia</i> | 1 | | | | | | | | X |
| Black Crane | <i>Amauromis flavirostris</i> | 2 | | | | | X | | X | |
| Common Moorhen | <i>Gallinula chloropus</i> | 2 | X | | | | X | | | |
| Red-knobbed Coot | <i>Fulica cristata</i> | 4 | X | X | X | | | | | X |
| Denham's Bustard | <i>Neotis denhami</i> | 1 | | X | | | | | | |
| Red-crested Korhaan | <i>Eupodotis ruficristata</i> | 1 | | | | X | | | | |
| African Jacana | <i>Actophilornis africanus</i> | 2 | | | | | X | | X | |
| Three-banded Plover | <i>Charadrius tricollaris</i> | 4 | | | | | X | X | X | X |
| Crowned Lapwing | <i>Vanellus coronatus</i> | 6 | X | | | X | X | X | X | X |
| Senegal Lapwing | <i>Vanellus lugubris</i> | 1 | | | | | | | X | |
| Black-winged Lapwing | <i>Vanellus melanopterus</i> | 1 | | X | | | | | | |
| Blacksmith Lapwing | <i>Vanellus armatus</i> | 8 | X | X | X | X | X | X | X | X |
| African Wattled Lapwing | <i>Vanellus senegallus</i> | 2 | X | X | | | | | | |
| Common Sandpiper | <i>Actitis hypoleucos</i> | 3 | | | | | X | X | X | |
| Wood Sandpiper | <i>Tringa glareola</i> | 2 | | | | | X | X | | |
| Common Greenshank | <i>Tringa nebularia</i> | 1 | | | | | | | | X |
| Ruff | <i>Philomachus pugnax</i> | 1 | | | | | | | | X |
| Black-winged Stilt | <i>Himantopus himantopus</i> | 3 | | | | | X | X | | X |
| Spotted Thick-knee | <i>Burhinus capensis</i> | 1 | | | | | | X | | |

| Common name | Scientific name | Total | Day | | | | | | | |
|----------------------------|----------------------------------|-------|-----|---|---|---|---|---|---|---|
| | | | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 |
| Water Thick-Knee | <i>Burhinus vermiculatus</i> | 3 | | | | | X | X | X | |
| Grey-headed Gull | <i>Larus cirrocephalus</i> | 2 | X | | | | | | | X |
| Whiskered Tern | <i>Chlidonias hybridus</i> | 2 | X | | | | | | | X |
| White-winged Tern | <i>Chlidonias leucopterus</i> | 1 | | | | | | | | X |
| Rock Dove | <i>Columba livia</i> | 3 | X | | X | X | | | | |
| Speckled Pigeon | <i>Columba guinea</i> | 2 | X | X | | | | | | |
| African Olive-Pigeon | <i>Columba arquatrix</i> | 3 | X | X | X | | | | | |
| Red-eyed Dove | <i>Streptopelia semitorquata</i> | 8 | X | X | X | X | X | X | X | X |
| African Mourning Dove | <i>Streptopelia decipiens</i> | 2 | | | | | X | X | | |
| Cape Turtle-Dove | <i>Streptopelia capicola</i> | 8 | X | X | X | X | X | X | X | X |
| Laughing Dove | <i>Streptopelia senegalensis</i> | 8 | X | X | X | X | X | X | X | X |
| Namaqua Dove | <i>Oena capensis</i> | 1 | | | | | X | | | |
| Emerald-spotted Wood-Dove | <i>Turtur chalcospilos</i> | 4 | | | | X | X | X | X | |
| Tambourine Dove | <i>Turtur tympanistria</i> | 1 | | | | | | | | X |
| African Green-Pigeon | <i>Treron calva</i> | 4 | | | | | X | X | X | X |
| Brown-headed Parrot | <i>Poicephalus cryptoxanthus</i> | 2 | | | | | X | X | | |
| Knysna Turaco (E) | <i>Tauraco corythaix</i> | 1 | | | X | | | | | |
| Purple-crested Turaco | <i>Tauraco porphyreolophus</i> | 2 | | | | | | | X | X |
| Grey Go-away Bird | <i>Corythaixoides concolor</i> | 5 | | | | X | X | X | X | X |
| Common Cuckoo | <i>Cuculus canorus</i> | 1 | | X | | | | | | |
| Red-chested Cuckoo | <i>Cuculus solitarius</i> | 1 | | | | X | | | | |
| Levaillant's Cuckoo | <i>Clamator levaillantii</i> | 3 | | | | X | X | | X | |
| Jacobin Cuckoo | <i>Clamator jacobinus</i> | 2 | | | | | X | X | | |
| Diderick Cuckoo | <i>Chrysococcyx caprius</i> | 4 | | X | | | X | X | X | |
| Burchell's Coucal (NE) | <i>Centropus burchellii</i> | 2 | | | | | X | X | | |
| Pearl-spotted Owllet | <i>Glaucidium perlatum</i> | 1 | | | | | | | X | |
| Cape Eagle-Owl | <i>Bubo capensis</i> | 1 | X | | | | | | | |
| Spotted Eagle-Owl | <i>Bubo africanus</i> | 1 | | | | | | X | | |
| European Nightjar | <i>Caprimulgus europaeus</i> | 1 | | | | | | X | | |
| Common Swift | <i>Apus apus</i> | 3 | | X | | | | | X | X |
| African Black Swift | <i>Apus barbatus</i> | 2 | | X | X | | | | | |
| White-rumped Swift | <i>Apus caffer</i> | 7 | X | X | X | X | | X | X | X |
| Little Swift | <i>Apus affinis</i> | 8 | X | X | X | X | X | X | X | X |
| Alpine Swift | <i>Apus melba</i> | 4 | X | X | X | | | | X | |
| African Palm Swift | <i>Cypsiurus parvus</i> | 5 | X | | X | | | X | X | X |
| Speckled Mousebird | <i>Colius striatus</i> | 6 | X | X | X | X | | X | X | |
| Red-faced Mousebird | <i>Urocolius indicus</i> | 5 | | | X | X | X | X | X | |
| Narina Trogon | <i>Apaloderma narina</i> | 1 | | | X | | | | | |
| Pied Kingfisher | <i>Ceryle rudis</i> | 4 | | | | | X | X | X | X |
| Giant Kingfisher | <i>Ceryle maxima</i> | 2 | | | | | X | X | | |
| Malachite Kingfisher | <i>Alcedo cristata</i> | 2 | | | | | X | | X | |
| Woodland Kingfisher | <i>Halcyon senegalensis</i> | 5 | | | | X | X | X | X | X |
| Brown-hooded Kingfisher | <i>Halcyon albiventris</i> | 4 | | | | X | X | | X | X |
| Striped Kingfisher | <i>Halcyon chelicuti</i> | 1 | | | | | | | X | |
| European Bee-eater | <i>Merops apiaster</i> | 5 | | | | X | X | X | X | X |
| Southern Carmine Bee-eater | <i>Merops nubicoides</i> | 4 | | | | X | X | X | X | |
| White-fronted Bee-eater | <i>Merops bullockoides</i> | 2 | | | | | | X | X | |

| Common name | Scientific name | Total | Day | | | | | | | |
|---------------------------------|---------------------------------|-------|-----|---|---|---|---|---|---|---|
| | | | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 |
| Little Bee-eater | <i>Merops pusillus</i> | 3 | | | | | X | X | X | |
| European Roller | <i>Coracias garrulus</i> | 5 | | | | X | X | X | X | X |
| Lilac-breasted Roller | <i>Coracias caudata</i> | 4 | | | | X | X | X | X | |
| Purple Roller | <i>Coracias naevia</i> | 1 | | | | | X | | | |
| African Hoopoe | <i>Upupa africana</i> | 4 | | | | X | X | X | X | |
| Green Wood-Hoopoe | <i>Phoeniculus purpureus</i> | 2 | | | | | | X | X | |
| Common Scimitarbill | <i>Rhinopomastus cyanomelas</i> | 2 | | | | | X | | X | |
| African Grey Hornbill | <i>Tockus nasutus</i> | 4 | | | | X | X | X | X | |
| Red-billed Hornbill | <i>Tockus erythrorhynchus</i> | 4 | | | | X | X | X | X | |
| Southern Yellow-billed Hornbill | <i>Tockus leucomelas</i> | 4 | | | | X | X | X | X | |
| Black-collared Barbet | <i>Lybius torquatus</i> | 4 | | | | X | | X | X | X |
| Yellow-fronted Tinkerbird | <i>Pogoniulus chrysoconus</i> | 1 | | | | X | | | | |
| Yellow-rumped Tinkerbird | <i>Pogoniulus bilineatus</i> | 1 | | | | X | | | | |
| Crested Barbet | <i>Trachyphonus vaillanti</i> | 3 | | | | | X | X | X | |
| Bennett's Woodpecker | <i>Campethera bennettii</i> | 2 | | | | | X | X | | |
| Golden-tailed Woodpecker | <i>Campethera abingoni</i> | 2 | | | | | | X | X | |
| Cardinal Woodpecker | <i>Dendropicos fuscescens</i> | 1 | | | | | X | | | |
| Bearded Woodpecker | <i>Dendropicos namaquus</i> | 2 | | | | | X | X | | |
| Rufous-naped Lark | <i>Mirafr africana</i> | 2 | | | | X | | X | | |
| Flappet Lark | <i>Mirafr rufocinnamomea</i> | 2 | | | | | | X | X | |
| Monotonous Lark | <i>Mirafr passeirina</i> | 2 | | | | | X | X | | |
| Sabota Lark (NE) | <i>Mirafr sabota</i> | 3 | | | | X | X | X | | |
| Eastern Long-billed Lark (E) | <i>Certhilauda semitorquata</i> | 1 | | X | | | | | | |
| Chestnut-backed Sparrowlark | <i>Eremopternix leucotis</i> | 1 | | | | | X | | | |
| Barn Swallow | <i>Hirundo rustica</i> | 8 | X | X | X | X | X | X | X | X |
| White-throated Swallow | <i>Hirundo albigularis</i> | 3 | X | X | X | | | | | |
| Wire-tailed Swallow | <i>Hirundo smithii</i> | 2 | | | | | | X | X | |
| Red-breasted Swallow | <i>Hirundo semirufa</i> | 3 | | | | X | X | X | | |
| Mosque Swallow | <i>Hirundo senegalensis</i> | 1 | | | | | | X | | |
| Greater Striped Swallow (E) | <i>Hirundo cucullata</i> | 4 | X | X | X | X | | | | |
| Lesser Striped Swallow | <i>Hirundo abyssinica</i> | 5 | | | X | X | | X | X | X |
| South African Cliff-Swallow (E) | <i>Hirundo spilodera</i> | 1 | | | | | | | | X |
| Rock Martin | <i>Hirundo fuligula</i> | 3 | | X | X | X | | | | |
| Common House Martin | <i>Delichon urbica</i> | 6 | | X | X | X | X | | X | X |
| Brown-throated Martin | <i>Riparia paludicola</i> | 1 | X | | | | | | | |
| Banded Martin | <i>Riparia cincta</i> | 2 | X | X | | | | | | |
| Black Saw-wing | <i>Psalidoprocne holomelas</i> | 1 | | | X | | | | | |
| Black Cuckooshrike | <i>Campephaga flava</i> | 2 | | | | | X | | | X |
| Fork-tailed Drongo | <i>Dicrurus adsimilis</i> | 7 | | X | X | X | X | X | X | X |
| Eurasian Golden Oriole | <i>Oriolus oriolus</i> | 2 | | | | | X | X | | |
| Black-headed Oriole | <i>Oriolus larvatus</i> | 4 | | X | | | | X | X | X |
| Cape Crow | <i>Corvus capensis</i> | 3 | X | X | X | | | | | |
| Pied Crow | <i>Corvus albus</i> | 3 | X | | X | X | | | | |
| Southern Black Tit | <i>Parus niger</i> | 5 | | | | X | X | X | X | X |
| Grey Penduline Tit | <i>Anthoscopus caroli</i> | 1 | | | | | X | | | |
| Arrow-marked Babbler | <i>Turdoides jardineii</i> | 5 | | | | X | X | X | X | X |
| Dark-capped Bulbul | <i>Pycnonotus barbatus</i> | 8 | X | X | X | X | X | X | X | X |

| Common name | Scientific name | Total | Day | | | | | | | |
|----------------------------------|-------------------------------------|-------|-----|---|---|---|---|---|---|---|
| | | | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 |
| Terrestrial Brownbul | <i>Phyllastrephus terrestris</i> | 2 | | | | X | | | X | |
| Yellow-streaked Greenbul | <i>Phyllastrephus flavostriatus</i> | 1 | | | X | | | | | |
| Sombre Greenbul | <i>Andropadus importunus</i> | 2 | | | | X | | | X | |
| Groundscraper Thrush | <i>Turdus litsitsirupa</i> | 2 | X | | | | | X | | |
| Cape Rock-Thrush (E) | <i>Monticola rupestris</i> | 2 | | | X | X | | | | |
| Sentinel Rock-Thrush (E) | <i>Monticola explorator</i> | 1 | | X | | | | | | |
| Mountain Wheatear (Chat) (NE) | <i>Oenanthe monticola</i> | 1 | | X | | | | | | |
| Buff-streaked Chat (E) | <i>Oenanthe bifasciata</i> | 1 | | X | | | | | | |
| Familiar Chat | <i>Cercomela familiaris</i> | 2 | | | X | X | | | | |
| Mocking Cliff-Chat | <i>Thamnolaea cinnamomeiventris</i> | 2 | | | X | X | | | | |
| Ant-eating Chat (E) | <i>Mymecocichla formicivora</i> | 3 | X | X | X | | | | | |
| African Stonechat | <i>Saxicola torquata</i> | 5 | X | X | X | X | | | | X |
| Chorister Robin-Chat (E) | <i>Cossypha dichroa</i> | 1 | | | X | | | | | |
| White-browed Robin-Chat | <i>Cossypha heuglini</i> | 1 | | | | | | | X | |
| Red-capped Robin-Chat | <i>Cossypha natalensis</i> | 1 | | | | X | | | | |
| Cape Robin-Chat | <i>Cossypha caffra</i> | 3 | | X | X | X | | | | |
| White-throated Robin-Chat (E) | <i>Cossypha humeralis</i> | 3 | | | | X | | X | X | |
| White-browed Scrub-Robin | <i>Erythropygia leucophrys</i> | 4 | | | | X | X | X | X | |
| Icterine Warbler | <i>Hippolais icterina</i> | 2 | | | | X | X | | | |
| Olive-tree Warbler | <i>Hippolais olivetorum</i> | 1 | | | | | X | | | |
| African Reed-Warbler | <i>Acrocephalus baeticatus</i> | 1 | | | | | | | X | |
| Marsh-Warbler | <i>Acrocephalus palustris</i> | 2 | | | | | | X | | X |
| Lesser Swamp-Warbler | <i>Acrocephalus gracilirostris</i> | 1 | | | | | | | X | |
| Dark-capped Yellow Warbler | <i>Chloropeta natalensis</i> | 1 | X | | | | | | | |
| Little Rush-Warbler | <i>Bradypterus baboecala</i> | 1 | | X | | | | | | |
| Willow Warbler | <i>Phylloscopus trochilus</i> | 5 | | | X | X | X | | X | X |
| Yellow-throated Woodland-Warbler | <i>Phylloscopus ruficapillus</i> | 1 | | | X | | | | | |
| Bar-throated Apalis | <i>Apalis thoracica</i> | 2 | | | X | X | | | | |
| Yellow-breasted Apalis | <i>Apalis flavida</i> | 2 | | | | X | | | X | |
| Long-billed Crombec | <i>Sylvietta rufescens</i> | 4 | | | | X | X | X | X | |
| Yellow-bellied Eremomela | <i>Eremomela icteropygialis</i> | 1 | | | | | X | | | |
| Green-capped Eremomela | <i>Eremomela scotops</i> | 1 | | | | | | | X | |
| Burnt-necked Eremomela | <i>Eremomela usticollis</i> | 2 | | | | | X | | X | |
| Green-backed Camaroptera | <i>Camaroptera brachyura</i> | 2 | | | | X | | | X | |
| Cape Grassbird (E) | <i>Sphenoecus afer</i> | 1 | | | X | | | | | |
| Zitting Cisticola | <i>Cisticola juncidis</i> | 6 | | X | X | X | | X | X | X |
| Desert Cisticola | <i>Cisticola aridulus</i> | 1 | | | | | | X | | |
| Wing-snapping Cisticola | <i>Cisticola ayresii</i> | 2 | | X | X | | | | | |
| Wailing Cisticola | <i>Cisticola lais</i> | 3 | | X | X | X | | | | |
| Rattling Cisticola | <i>Cisticola chiniana</i> | 5 | | | | X | X | X | X | X |
| Red-faced Cisticola | <i>Cisticola erythrops</i> | 2 | | | | | | X | X | |
| Levaillant's Cisticola | <i>Cisticola tinniens</i> | 2 | X | X | | | | | | |
| Croaking Cisticola | <i>Cisticola natalensis</i> | 1 | | | | | | | | X |
| Lazy Cisticola | <i>Cisticola aberrans</i> | 1 | | X | | | | | | |
| Neddicky | <i>Cisticola fulvicapillus</i> | 1 | | | | | | | X | |
| Tawny-flanked Prinia | <i>Prinia subflava</i> | 4 | | | | | X | X | X | X |

| Common name | Scientific name | Total | Day | | | | | | | |
|----------------------------------|-----------------------------------|-------|-----|---|---|---|---|---|---|---|
| | | | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 |
| Drakensberg Prinia (E) | <i>Prinia hypoxantha</i> | 4 | X | X | X | X | | | | |
| Spotted Flycatcher | <i>Muscicapa striata</i> | 6 | | | X | X | X | X | X | X |
| African Dusky Flycatcher | <i>Muscicapa adusta</i> | 1 | | | X | | | | | |
| Ashy Flycatcher | <i>Muscicapa caerulescens</i> | 1 | | | | | | | X | |
| Grey Tit-Flycatcher | <i>Myioparus plumbeus</i> | 3 | | | | | X | X | X | |
| Southern Black Flycatcher | <i>Melaenornis pammelaina</i> | 3 | | | | | | X | X | X |
| Cape Batis (E) | <i>Batis capensis</i> | 1 | | | X | | | | | |
| Chin-spot Batis | <i>Batis molitor</i> | 5 | | | | X | X | X | X | X |
| Blue-mantled Crested-Flycatcher | <i>Trochocecus cyanomelas</i> | 1 | | | X | | | | | |
| African Paradise Flycatcher | <i>Terpsiphone viridis</i> | 4 | | X | X | X | | | X | |
| African Pied Wagtail | <i>Motacilla aguimp</i> | 3 | | | | | X | X | X | |
| Mountain Wagtail | <i>Motacilla clara</i> | 2 | | | X | X | | | | |
| Cape Wagtail | <i>Motacilla capensis</i> | 4 | X | X | X | X | | | | |
| African Pipit | <i>Anthus cinnamomeus</i> | 2 | | X | | | X | | | |
| Long-billed Pipit | <i>Anthus similis</i> | 1 | | X | | | | | | |
| Striped Pipit | <i>Anthus lineiventris</i> | 1 | | | | X | | | | |
| Bushveld Pipit | <i>Anthus caffer</i> | 1 | | | | | | | X | |
| Yellow-breasted Pipit (E) | <i>Hemimacronyx chloris</i> | 1 | | X | | | | | | |
| Cape Longclaw (E) | <i>Macronyx capensis</i> | 2 | | X | X | | | | | |
| Yellow-throated Longclaw | <i>Macronyx croceus</i> | 3 | | | | X | | X | X | |
| Lesser Grey Shrike | <i>Lanius minor</i> | 2 | | | | | X | X | | |
| Common Fiscal | <i>Lanius collaris</i> | 5 | X | X | X | X | | | | X |
| Red-backed Shrike | <i>Lanius collurio</i> | 5 | | | | X | X | X | X | X |
| Magpie Shrike | <i>Corvinella melanoleuca</i> | 4 | | | | X | X | X | X | |
| Southern Boubou (E) | <i>Laniarius ferrugineus</i> | 2 | | | | X | | X | | |
| Black-backed Puffback | <i>Dryoscopus cubla</i> | 5 | | | | X | X | X | X | X |
| Brubru | <i>Nilaus afer</i> | 1 | | | | | X | | | |
| Brown-crowned Tchagra | <i>Tchagra australis</i> | 1 | | | | | | | X | |
| Black-crowned Tchagra | <i>Tchagra senegala</i> | 2 | | | | | | X | X | |
| Bokmakierie (E) | <i>Telophorus zeylonus</i> | 1 | | X | | | | | | |
| Orange-breasted Bush-Shrike | <i>Telophorus sulfureopectus</i> | 1 | | | | | | X | | |
| Grey-headed Bush-Shrike | <i>Malaconotus blanchoti</i> | 1 | | | | | | X | | |
| White-crested Helmet-Shrike | <i>Prionops plumatus</i> | 2 | | | | | | X | X | |
| Southern White-crowned Shrike | <i>Eurocephalus anguitimens</i> | 1 | | | | | X | | | |
| Common Myna | <i>Acridotheres tristis</i> | 5 | X | X | X | X | | | | X |
| Pied Starling (E) | <i>Spreo bicolor</i> | 4 | X | X | X | | | | | X |
| Wattled Starling | <i>Creatophora cinerea</i> | 4 | X | | | X | X | X | | |
| Violet-backed Starling | <i>Cinnyricinclus leucogaster</i> | 4 | | | | | X | X | X | X |
| Burchell's Starling (NE) | <i>Lamprotomis australis</i> | 4 | | | | X | X | X | X | |
| Cape Glossy Starling | <i>Lamprotomis nitens</i> | 2 | | | | | | | X | X |
| Greater Blue-eared Starling | <i>Lamprotomis chalybaeus</i> | 3 | | | | | X | X | X | |
| Red-winged Starling | <i>Onychognathus morio</i> | 5 | X | X | X | X | X | | | |
| Red-billed Oxpecker | <i>Buphagus erythrorhynchus</i> | 4 | | | | X | X | X | X | |
| Gurney's Sugarbird (E) | <i>Promerops gurneyi</i> | 1 | | X | | | | | | |
| Malachite Sunbird | <i>Nectarinia famosa</i> | 4 | X | X | X | X | | | | |
| Sthn Double-collared Sunbird (E) | <i>Nectarinia chalybea</i> | 1 | | | X | | | | | |

| Common name | Scientific name | Total | Day | | | | | | | |
|---------------------------------|---------------------------------|------------|-----------|-----------|-----------|------------|------------|------------|------------|------------|
| | | | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 |
| Gtr Double-collared Sunbird (E) | <i>Nectarinia afra</i> | 2 | | | X | X | | | | |
| White-bellied Sunbird | <i>Nectarinia talatala</i> | 2 | | | | X | | | | X |
| Scarlet-chested Sunbird | <i>Nectarinia senegalensis</i> | 2 | | | | | | | X | X |
| Amethyst Black Sunbird | <i>Nectarinia amethystina</i> | 2 | | X | X | | | | | |
| Collared Sunbird | <i>Anthreptes collaris</i> | 2 | | | | | | X | X | |
| Cape White-eye (E) | <i>Zosterops pallidus</i> | 6 | X | X | X | X | | | X | X |
| Red-billed Buffalo-Weaver | <i>Bubalomis niger</i> | 4 | | | | X | X | X | X | |
| House Sparrow | <i>Passer domesticus</i> | 6 | X | | X | X | X | | X | X |
| Cape Sparrow (NE) | <i>Passer melanurus</i> | 1 | X | | | | | | | |
| Southern Grey-headed Sparrow | <i>Passer diffusus</i> | 5 | | | | X | X | X | X | X |
| Yellow-throated Petronia | <i>Petronia supercilii</i> | 1 | | | | | | X | | |
| Thick-billed Weaver | <i>Amblyospiza albifrons</i> | 1 | | | | X | | | | |
| Spectacled Weaver | <i>Ploceus ocularis</i> | 1 | | | | | | | X | |
| Village Weaver | <i>Ploceus cucullatus</i> | 1 | | | | | X | | | |
| Cape Weaver (E) | <i>Ploceus capensis</i> | 2 | X | X | | | | | | |
| Southern Masked-Weaver | <i>Ploceus velatus</i> | 6 | X | X | X | X | X | X | | |
| Lesser Masked-Weaver | <i>Ploceus intermedius</i> | 2 | | | | | | X | X | |
| Red-billed Quelea | <i>Quelea quelea</i> | 2 | | | | | X | X | | |
| Southern Red Bishop | <i>Euplectes orix</i> | 6 | X | X | X | | X | | X | X |
| Yellow-crowned Bishop | <i>Euplectes afer</i> | 1 | X | | | | | | | |
| Yellow Bishop | <i>Euplectes capensis</i> | 1 | | X | | | | | | |
| Fan-tailed Widowbird | <i>Euplectes axillaris</i> | 4 | X | X | X | | | | | X |
| White-winged Widowbird | <i>Euplectes albonotatus</i> | 4 | | | X | X | X | X | | |
| Red-collared Widowbird | <i>Euplectes ardens</i> | 4 | | X | X | | | | X | X |
| Long-tailed Widowbird | <i>Euplectes progne</i> | 4 | X | X | X | | | | | X |
| Green-winged Pytilia | <i>Pytilia melba</i> | 3 | | | | X | X | X | | |
| Jameson's Firefinch | <i>Lagonosticta rhodopareia</i> | 2 | | | | X | | X | | |
| Red-billed Firefinch | <i>Lagonosticta senegala</i> | 2 | | | | | | X | X | |
| Blue Waxbill | <i>Uraeginthus angolensis</i> | 5 | | | | X | X | X | X | X |
| Common Waxbill | <i>Estrilda astrild</i> | 5 | X | X | X | | X | | X | |
| Orange-breasted Waxbill | <i>Sporaeginthus subflavus</i> | 1 | X | | | | | | | |
| Red-headed Finch (NE) | <i>Amadina erythrocephala</i> | 1 | X | | | | | | | |
| Pin-tailed Whydah | <i>Vidua macroura</i> | 6 | X | X | X | | | X | X | X |
| Long-tailed Paradise-Whydah | <i>Vidua paradisaea</i> | 4 | | | | X | X | X | X | |
| Purple Indigobird | <i>Vidua purpurascens</i> | 1 | | | | X | | | | |
| Village Indigobird | <i>Vidua chalybeata</i> | 3 | | | | | X | X | X | |
| Yellow-fronted Canary | <i>Serinus mozambicus</i> | 5 | | | | X | X | X | X | X |
| Cape Canary | <i>Serinus canicollis</i> | 3 | X | X | X | | | | | |
| Streaky-headed Seed-Eater | <i>Serinus gularis</i> | 1 | | | | X | | | | |
| Golden-breasted Bunting | <i>Emberiza flaviventris</i> | 3 | | | | | X | | X | X |
| Cinnamon-breasted Bunting | <i>Emberiza tahapisi</i> | 3 | | | | X | X | | | X |
| TOTAL | 319 | 890 | 82 | 88 | 91 | 122 | 133 | 139 | 135 | 100 |

Mammal List

| Common name | Scientific name | Total | Day | | | | | | | | |
|------------------------------|------------------------------------|-----------|-----------|----------|----------|----------|-----------|-----------|-----------|-----------|----------|
| | | | 1 | 2 | 3 | 4 | 5 | 6 | 7 | 8 | |
| Peter's Epauletted Fruit Bat | <i>Epomophorus crypturus</i> | 1 | | | | | | | | X | |
| Tomb Bat | <i>Taphozous mauritanus</i> | 1 | | | | | | | X | | |
| Warthog | <i>Phacochoerus aethiopicus</i> | 5 | | | | X | X | X | X | X | |
| White Rhinoceros | <i>Ceratotherium simum</i> | 2 | | | | | | X | X | | |
| Hippopotamus | <i>Hippopotamus amphibius</i> | 3 | | | | | X | X | X | | |
| Southern Giraffe | <i>Giraffa camelopardalis</i> | 4 | | | | X | X | X | X | | |
| Common Duiker | <i>Sylvicapra grimmia</i> | 2 | | X | | | | X | | | |
| Eland | <i>Taurotragus oryx</i> | 1 | | | X | | | | | | |
| Bushbuck | <i>Tragelaphus scriptus</i> | 3 | | | | | | X | X | X | |
| Greater Kudu | <i>Tragelaphus strepsiceros</i> | 3 | | | | | | X | X | X | |
| Common Reedbuck | <i>Redunca arundinum</i> | 1 | | X | | | | | | | |
| Mountain Reedbuck | <i>Redunca fulvorufula</i> | 1 | X | | | | | | | | |
| Common Waterbuck | <i>Kobus ellipsiprymnus</i> | 4 | | | | X | X | X | X | | |
| Impala | <i>Aepyceros melampus</i> | 5 | | | | X | X | X | X | X | |
| Klipspringer | <i>Oreotragus oreotragus</i> | 3 | | | X | | | | | X | X |
| Steenbok | <i>Raphicerus campestris</i> | 4 | X | | | X | | | X | X | |
| Oribi | <i>Ourebia ourebi</i> | 1 | | X | | | | | | | |
| Grey Rhebok | <i>Palea capreolus</i> | 2 | | X | X | | | | | | |
| Blesbok | <i>Damaliscus dorcas phillipsi</i> | 2 | | X | X | | | | | | |
| Blue Wildebeest | <i>Connochaetes taurinus</i> | 4 | | | | X | X | X | X | | |
| Cape Buffalo | <i>Syncerus caffer</i> | 2 | | | | | X | X | | | |
| Burchell's Zebra | <i>Equus burchelli</i> | 4 | | | | X | X | X | X | | |
| African Elephant | <i>Loxodonta africana</i> | 4 | | | | X | X | X | X | | |
| Greater Cane Rat | <i>Thryonomys swinderianus</i> | 1 | X | | | | | | | | |
| Bushveld Gerbil | <i>Tatera leucogaster</i> | 1 | | | | | | X | | | |
| Tree Squirrel | <i>Paraxerus cepapi</i> | 5 | | | | X | X | X | X | X | |
| Scrub Hare | <i>Lepus saxatilis</i> | 1 | | | | | | X | | | |
| Black-backed Jackal | <i>Canis mesomelas</i> | 2 | | X | | X | | | | | |
| Suricate | <i>Suricata suricatta</i> | 1 | X | | | | | | | | |
| Banded Mongoose | <i>Mungos mungo</i> | 1 | | | | | | | | X | |
| Dwarf Mongoose | <i>Helogale parvula</i> | 3 | | | | | X | | X | X | |
| Slender Mongoose | <i>Galerella sanguinea</i> | 4 | | | | X | X | X | X | | |
| Yellow Mongoose | <i>Cynictis penicillata</i> | 1 | | X | | | | | | | |
| Spotted Hyaena | <i>Crocuta crocuta</i> | 1 | | | | | | X | | | |
| Lion | <i>Panthera leo</i> | 1 | | | | | | | | | X |
| Leopard | <i>Panthera pardus</i> | 2 | | | | X | X | | | | |
| Chacma Baboon | <i>Papio ursinus</i> | 5 | | | X | X | X | X | X | | |
| Vervet Monkey | <i>Cercopithecus aethiops</i> | 7 | | X | X | X | X | X | X | X | X |
| Samango Monkey | <i>Cercopithecus mitis</i> | 1 | | | X | | | | | | |
| TOTAL | | 39 | 99 | 4 | 8 | 7 | 14 | 18 | 21 | 20 | 7 |